Granudacyn® irrigation solution and gel for cleaning, moistening and rinsing

Granudacyn is a wound irrigation solution for cleansing and moisturising acute, chronic and contaminated wounds as well as for superficial and partial thickness burns. Hypochlorous acid (HOCl) ensures safe preservation and makes Granudacyn a reliable wound irrigation solution. HOCl prevents the proliferation of Gram+ and Gram- bacteria including; MRSA, ORSA, VRSA, VRE, viruses, fungi and spores¹.

Granudacyn advantages

- Cleans the wound mechanically
- Is free of microorganisms and pH neutral
- Is non-cytotoxic and non-irritant
- Is hypotonic
- Is free of heavy metals
- Reduces wound malodour²

- Shelf life 24 months for irrigation solution and 18 months for gel
- First choice for peritoneal lavage²
- Can be applied on CNS tissue, cartilage and bones
- Well suited for cavities and fistulas
- · Can remain in the wound





How Granudacyn® works

Granudacyn is preserved to allow multi-patient use for up to 60 days (solution) and 90 days (gel) after opening. To ensure the safe use after opening, the Granudacyn products are preserved with hypochlorous acid, a substance naturally generated in the human body by activated neutrophils².




1. Disrupts the cell wall structures Granudacyn Irrigation solution surrounds the microorganisms, and the hypochlorus acid penetrates the cell wall of the microorganism and increases its permeability³.



2. Osmolysis leads to cell burst. The hypotonic nature of the Granudacyn Irrigation solution causes water to flow into the cells. The result is osmolysis: the increasing internal pressure causes the cells to burst.

Cytotoxicity4

Granudacyn caused neither toxicological nor biological damages to a subconfluent layer of mousefibroblasts (L929). In contrast, the test substance with a NaOCl concentration of 800 ppm led to cytotoxic reaction in mouse fibroblasts³. Granudacyn is graded not cytotoxic.



Composition: Granudacyn contains water, sodium chlorite, sodium hypochlorite and hypochlorous acid. The relative concentration of HOCl and NaOCl is pH dependent. At neutral pH Granudacyn Wound Irrigation solution contains 50 ppm HOCl and 50 ppm NaOCl and Wound gel contains 40 ppm HOCl and 40 ppm NaOCl.

Storage: Granudacyn solution has a shelf life of 24 months after manufacturing and 60 days after opening. Granudacyn wound gel has a shelf life of 18 months after manufacturing and 90 days after opening. Granudacyn solution and gel can be stored at room temperature.

Product information

Product	Size	Product code	Shelf life	Pcs/box
Granudacyn Wound irrigation solution	50ml spray	360150	24 months	20
	250ml spray	360100		15
	500ml	360101		12
	1000ml	360102		6
	500ml NPWT	360103		12
	1000ml NPWT	360104		6
Granudacyn Wound gel	50g	360107	18 months	12
	100g spray	360108		12
	250g spray	360106		15

Granudacyn can be used for the irrigation, cleaning and decontamination of the following wound-types:

- All chronic wounds of any depths, such as diabetic foot ulcers, pressure injuries, venous leg ulcers, etc.
- All acute wounds, such as cuts, bite wounds, lacerations, abrasions, etc.
- Surgical wounds (intraoperative and postoperative)
- Wounds with exposed cartilage, tendons, ligaments and/or bones
- Superficial and partial thickness burns
- Radiation ulcers
- Fistulas and abscesses

References

- 1. In-vitro suspension test (EN13727, EN 13624, EN 13704, EN 14476 phase 2) with Granudacyn® wound irrigation solution.
- 2. Consensus on Wound Antisepsis: Update 2018, Skin Pharmacol Physiol 2018;31:28-58, DOI: 10.1159/000481545.
- 3. Fukuzaki, Biocontrol Science, 2006, Vol. 11, No. 4, 147-157.
- $4.\ Method\ Ph. Eur.\ 2.2.35, test\ conducted\ by\ BIOSERV\ Analytik-\ und\ Medizin produkte\ GmbH,\ Rostock,\ Germany.$

www.molnlycke.com.au | www.molnlycke.co.nz

